

*Pieris brassicae* L.

This species would seem to be more numerous than I perhaps indicated, as Mr. H. G. Allcard tells me that he found it commonly in the vegetable growing area inland of Palma. He also comments on its small size.

*Artogeia napi* L.

In the postscript to my paper I noted the inclusion of *A. napi* as a Majorcan species by Bustillo and Rubio (1974). Señor M. R. Gomez Bustillo has now informed me, however, that this has subsequently been found to be an error due to the incorrect identification of a single female which was in fact *A. rapae* L.

## References

- Bustillo, M. R. G. and Rubio, F. F., 1974. *Mariposas de la Peninsula Iberica*. Madrid.  
 Manley, W. B. L. and Allcard, H. G., 1970. *A Field Guide to the Butterflies and Burnets of Spain*.  
 Muschamp, P. A. H., 1904. Majorca — Eight Days Entomology, Two New Butterfly Aberrations. *Ent. Rec.*, **16**: 221-223.  
 Perceval, M. J., 1974. Butterflies of Majorca. *Ent. Rec.*, **86**: 225-234.

## Notes and Observations

EUDEMIS PORPHYRANA (HUEBNER) NEW TO KENT. — A single specimen of this Tortricid appeared in the m.v. trap in my garden here on the night of August 4th/5th, 1976. So far as I am aware this is the first record of this species for Kent. The moth appears to have been first recognised as British in 1915 from Devon and Herefordshire, whence it seems to have extended its range eastwards. — J. M. CHALMERS-HUNT.

APOMYELOIS BISTRIATELLA NEOPHANES (DURRANT, 1915) (LEP.: PYRALIDAE) IN WALES. — I would like, rather belatedly, to record breeding this moth in September 1974 from a larva that I found in *Daldinia concentrica* on *Ulex*, on sea cliffs near Oxwich, Gower Peninsula, Glamorgan. I have not heard of any other South Wales records for this species. — Dr. A. N. B. SIMPSON, 29 The Greenway, Colletts Green, Powick, near Worcester.

THE CONVULVULUS HAWK IN HANTS. — On 22nd September, 1976, I discovered a male *Agrius convolvuli* L. clinging to the bottom of a sheet hanging near the m.v. trap in my garden, and on the following morning a further male was found in almost exactly the same place. Later that evening, my son found yet another male on the ground about six feet from the trap. Apparently it had been there all day but had fortunately avoided the attentions of the birds. I was also given a dead *convolvuli* which had been found on the wall of a house at BurrIDGE, near Southampton, on 22nd September. — D. A. LEPARD, Silver Crest, Silver Street, Sway, Lymington, Hants.